

Losing control of your bladder may have nothing to do with your age!

It is possible to take control of your bladder and reduce your symptoms.

Know

About the different types of incontinence:

- Urgency incontinence you often have uncontrollable urges to urinate
- Stress incontinence you urinate when you sneeze, cough or laugh
- Functional incontinence you can't get to the bathroom fast enough
- Overflow incontinence your bladder is full and causes dribbling and difficulty with emptying your bladder

Ask

- A doctor or nurse about re-training your bladder
- Your pharmacist or doctor to look at the full list of medications you are taking to see if any of them are contributing to incontinence
- Your doctor about what kind of physical activity and how much is right for you. It's important to stay active!

Do

- Wear pants with minimal number of buttons or zippers so that fewer things get in the way of your urinating
- Use unscented, moistened disposable wipes, or a warm washcloth to clean up after an accident and change into clean clothes as soon as possible
- Pelvic floor exercises, as directed by a doctor, nurse, or physiotherapist, to strengthen the muscles that support your bladder and urethra
- Drink plenty of non-caffeinated beverages to stay hydrated. If you don't drink enough water, your bladder lining may become irritated and increase your incontinence symptoms
- Exercise regularly and eat a nutritious diet to maintain a healthy weight
- Encourage your older family member to use the toilet at regularly scheduled intervals (For caregivers - if your family member is the one living with incontinence as well as a cognitive impairment)



